

# magnesium sulfate

## 2.465g/5mL or 2.5g/5mL injection\*

© Department for Health and Ageing, Government of South Australia. All rights reserved.

**Note:**

This guideline provides advice of a general nature. This statewide guideline has been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach. The guideline is based on a review of published evidence and expert opinion.

Information in this statewide guideline is current at the time of publication.

SA Health does not accept responsibility for the quality or accuracy of material on websites linked from this site and does not sponsor, approve or endorse materials on such links.

Health practitioners in the South Australian public health sector are expected to review specific details of each patient and professionally assess the applicability of the relevant guideline to that clinical situation.

If for good clinical reasons, a decision is made to depart from the guideline, the responsible clinician must document in the patient's medical record, the decision made, by whom, and detailed reasons for the departure from the guideline.

This statewide guideline does not address all the elements of clinical practice and assumes that the individual clinicians are responsible for discussing care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes:

- The use of interpreter services where necessary,
- Advising consumers of their choice and ensuring informed consent is obtained,
- Providing care within scope of practice, meeting all legislative requirements and maintaining standards of professional conduct, and
- Documenting all care in accordance with mandatory and local requirements

**This is a High Risk Medication** 

An overdose can be fatal.

## Dose and Indications

\* There are two preparations available:

2.465g/5mL (49.3%) magnesium sulfate contains 10mmol magnesium

2.5g/5mL (50%) magnesium sulfate heptahydrate contains 10.3mmol magnesium

In dosing with neonates, the difference becomes immeasurable and for the purpose of these guidelines, they each contain elemental magnesium 10mmol/5mL or 2mmol/mL

All doses must be written as millimoles of elemental magnesium

## Hypomagnesaemia

### Intravenous, Intramuscular

0.2 to 0.4mmol/kg/dose every 12 hours as required

## Pulmonary Hypertension

### Intravenous

Loading dose 0.8mmol/kg then if clinical response continue

Maintenance dose 0.1 to 0.3mmol/kg/hr for 2 to 5 days

## Torsades de pointes (Prolonged QT syndrome due to hypomagnesaemia)

### Intravenous

0.1 to 0.2mmol/kg/dose

**INFORMAL COPY WHEN PRINTED**

Page 1 of 4



# magnesium sulfate

## 2.465g/5mL or 2.5g/5mL injection\*

### Preparation and Administration

#### Intravenous

Dilute 2mL of either product (2mmol/mL of elemental magnesium) with 3mL of compatible fluid (to give a total volume of 5mL). The resulting solution contains 0.8mmol/mL magnesium.

Dose	0.2mmol	0.4mmol	0.6mmol	0.8mmol	1mmol	1.2mmol
Volume	0.25mL	0.5mL	0.75mL	1mL	1.25mL	1.5mL

Further dilute with a compatible fluid.

Low doses can be given over 20 minutes.

Doses greater than 0.8mmol/kg should be given at a maximum recommended rate of 0.04mmol/kg/minute.

#### Intramuscular

Dilute 2mL of either product (2mmol/mL of elemental magnesium) with 3mL of compatible fluid (to give a total volume of 5mL). The resulting solution contains 0.8mmol/mL magnesium.

Dose	0.2mmol	0.4mmol	0.6mmol	0.8mmol	1mmol	1.2mmol
Volume	0.25mL	0.5mL	0.75mL	1mL	1.25mL	1.5mL

Intramuscular is painful and sometimes causes haematomas.

### Compatible Fluids

Glucose 5%, glucose/sodium solutions, sodium chloride 0.9%

### Adverse Effects

#### Common

Flushing, vomiting

Other adverse effects are often related to the development of hypermagnesaemia: important signs are loss of deep tendon reflexes and respiratory depression. More serious effects are hypotension, bradycardia, CNS depression, coma, circulatory collapse, cardiac arrest.



# magnesium sulfate

## 2.465g/5mL or 2.5g/5mL injection\*

---

### Monitoring

- > Magnesium levels regularly; usual blood level 0.75 to 1mmol/L however if treating pulmonary hypertension aim for levels between 3.5 to 5.5mmol/L
- > Renal function
- > Urine output
- > Electrolytes
- > Blood pressure, heart rate, respiratory rate, oxygen saturation, urine output, reflexes and other sign of toxicity regularly during treatment is recommended.

### Practice Points

- > All doses must be written as elemental magnesium
- > Anticipate changes in calcium and phosphorus balance
- > Calcium gluconate 10% injection should be available in case of hypermagnesaemia
- > Use CAUTIOUSLY in patients with renal impairment and/or electrolyte imbalance
- > DO NOT USE in patients with heart block or myocardial damage
- > When treating pulmonary hypertension consider other agents (nitric oxide and sildenafil) before using magnesium sulphate.



# magnesium sulfate

## 2.465g/5mL or 2.5g/5mL injection\*

### Document Ownership & History

<b>Developed by:</b>	SA Maternal, Neonatal & Gynaecology Community of Practice
<b>Contact:</b>	<a href="mailto:Health.NeoMed@sa.gov.au">Health.NeoMed@sa.gov.au</a>
<b>Endorsed by:</b>	SA Health Safety and Quality Strategic Governance Committee
<b>Next review due:</b>	31/05/2020
<b>ISBN number:</b>	978-1-74243-409-4
<b>PDS reference:</b>	<b>CG038</b>
<b>Policy history:</b>	Is this a new policy (V1)? <b>N</b> Does this policy amend or update an existing policy? <b>Y</b> If so, which version? <b>2.0</b> Does this policy replace another policy with a different title? <b>N</b> If so, which policy (title)?

Approval Date	Version	Who approved New/Revised Version	Reason for Change
9/03/18	V2.1	SA Health Safety and Quality Strategic Governance Committee	Review date extended to 5 years following risk assessment. New template.
03/2015	V2	SA Health Safety and Quality Strategic Governance Committee	High risk notification included
11/2012	V1	SA Maternal & Neonatal Clinical Network	Original SA Maternal & Neonatal Clinical Network approved version

